

Reducibility, Arrhenius plots and the Uroboros Dragon¹,
a reply to the preprint

“Correlations in Nuclear Arrhenius-Type Plots”

by M.B. Tsang and P. Danielewicz.

L.G. Moretto, L. Beaulieu, L. Phair, and G.J. Wozniak

*Nuclear Science Division, Lawrence Berkeley National
Laboratory, Berkeley, CA 94720, USA*

June 5, 1997

In previous publications [1–3], which have generated more attention than anticipated [4–9], we showed that the n -fragment multifragmentation probabilities at any given transverse energy $P_n(E_t)$, could be reduced by means of the binomial expression

$$P_n^m = \frac{m!}{n!(m-n)!} p^n (1-p)^{m-n} \quad (1)$$

to a single one fragment probability $p(E_t)$, where $E_t = \sum E_i \sin^2 \theta_i$, the sum of the kinetic energies of all charged particles in an event weighted by the sine squared of their polar angle. This “reducibility” with E_t was taken as a strong indication that fragments were emitted nearly independently of one another with the same probability $p(E_t)$. We further noticed that, under the assumption that E_t is proportional to the excitation energy E (and thus temperature $T \propto \sqrt{E_t}$), the Arrhenius plots of $\log(1/p)$ versus $1/\sqrt{E_t}$, produced straight lines. We called this “thermal scaling” because it suggests the Boltzmann form for the elementary probability $p \propto \exp(-B/T)$ or

$$\log \frac{1}{p} = \frac{B}{T} = \frac{B'}{\sqrt{E_t}}, \quad (2)$$

¹ The Uroboros dragon, or the snake eating its own tail, is an archetype that can be traced from the Egyptians through the Greeks and the Gnostic religions into the Alchemical writings of the Middle Ages and Renaissance all the way to modern times. A symbol of self reference (think of Escher’s hand drawing the hand that draws it...), it has been used to explain the universe. It has been applied by the authors of the preprint with a somewhat more modest intent.

with B representing the barrier for fragment emission.

The present preprint attacks both of our claims. The first on the basis that our binomial fits could be matched in quality by a constrained Poissonian function. The second on the basis that the Arrhenius plots are allegedly shown to be the result of a strong autocorrelation.

Since the second criticism is potentially more devastating than the first, we begin with it.

As a preemptive strike, the authors raise concerns about the relevance of using $\sqrt{E_t}$ as a measure of the temperature. They cite as evidence to the contrary recent work exploring the impact parameter dependence of temperatures extracted from excited state populations [10]. In that work a value of $T=4.6$ MeV is determined, largely independent of impact parameter for moderately central to very central collisions of Au+Au at 35 MeV/nucleon. We are not unduly concerned by this result since temperatures of 4-5 MeV are almost *always* extracted from such studies, independent of bombarding energy [11,12], entrance channel mass asymmetry [11,12], as well as impact parameter [10,13]. Since such extracted temperatures seem to be insensitive to *most* features of heavy-ion reactions, we would not be surprised to find that they are also insensitive to E_t .

Furthermore, temperatures extracted from the double isotope ratio method [14] do indeed show a broad range of temperatures for heavy ion reactions at intermediate energies [15].

Coming to the heart of their criticism, the authors argue that the observed linearity of the Arrhenius plots is due to autocorrelation. However, a fatal flaw mars the derivation of the proposed explanation for linear Arrhenius plots [9] and undermines the conclusions of the preprint. This flaw is circular logic in the algebra, leading to a trivial identity that always holds, regardless of whether or not there exists an autocorrelation.

Assuming with the authors that the binomial number of “throws” m is constant (for many experiments this is a good assumption) we consider the Arrhenius plot constructed with the average intermediate mass fragment (IMF) multiplicity $\langle n \rangle = mp$ and E_t . So we plot $\log 1/\langle n \rangle$ versus $1/\sqrt{E_t}$ and in all experimental cases we obtain a straight line. As we have observed [3], the authors also observe that E_t is constructed both with the light charged particles and the IMFs. In the authors’ notation

$$E_t = (N_C - \langle n \rangle) E_t^{\text{LP}} + \langle n \rangle E_t^{\text{IMF}}, \quad (3)$$

where N_C is the number of charged particles in an event and E_t^{LP} and E_t^{IMF} are




Fig. 1. Top panel: The transverse energy E_t as a function of the average IMF multiplicity $\langle n \rangle$ for the reaction Ar+Au, $E/A=110$ MeV. The line is a straight-line fit, Eq. (5). Bottom panel: $\langle n \rangle$ as a function of E_t . The fit is the exponential fit of Eq. (9) used in refs. [1–3].

the average transverse energies of the light particles and IMFs, respectively. The use of $\langle n \rangle$ to construct this independent variable E_t creates the danger of autocorrelation in the study of the dependence of $\langle n \rangle$ on E_t (as in the Arrhenius plot). This we also have discussed extensively [3].

Now the authors [9] observe that $\langle n \rangle$ and N_C are globally very strongly correlated. They fit this correlation with the linear form

$$N_C = a' + b' \langle n \rangle \quad (4)$$

and finally they express E_t as a function of the single variable $\langle n \rangle$

$$E_t = b \left(\frac{a}{b} + \langle n \rangle \right) \quad (5)$$

where $a = a' E_t^{\text{LP}}$ and $b = (b' - 1) E_t^{\text{LP}} + E_t^{\text{IMF}}$.

A quicker but nearly equivalent procedure would have been to fit the equally

strongly correlated quantities E_t , $\langle n \rangle$ directly with the above form (see the top panel of Fig. 1) since N_C and E_t are very tightly correlated as well. The values of a and b from a direct fit using Eq. (5) are 100 MeV and 220 MeV respectively. These values are nearly equal to those extracted by the authors ($a=106$ MeV, $b=213$ MeV) when starting with Eq. (4).

A moment's reflection shows that Eq. (5) contains the original experimental dependence of $\langle n \rangle$ on E_t (bottom panel of Fig. 1), and it does not differ in any significant way, other than in the linear fit, from the data used to construct the Arrhenius plots. It is just its inverse! Here is the start of the vicious circle.

In fact, to fit the data they could have used with better results the Boltzmann-like functional form which gives linear Arrhenius plots as we have done in the bottom panel of Fig. 1. One can argue on which of the two is a better fit, but not on the fact that, for the two functions, one is approximately the inverse of the other.

Armed with Eq. (5), the authors take an experimental Arrhenius plot which is linear, and they substitute E_t with the value calculated from Eq. (5). Since the linear form is a good approximation to E_t , nothing much happens to the Arrhenius plot which of course remains linear. It simply means that the form of Eq. (5) is a good approximation to the data!

Yet the authors make the surprising claim that the linearity in such a plot is the result of an autocorrelation because now one has $\langle n \rangle$ both in the ordinate (as $\log 1/\langle n \rangle$) and in the abscissa (as $1/\sqrt{b(a/b + \langle n \rangle)}$).

In what follows we show that this technique for demonstrating autocorrelations is so general and fundamentally flawed, that it will give the *same* result for all functions, *even for those in which autocorrelation has been ruled out by construction*. We call this method the Uroboros method, for reasons that will become clear below.

As we noticed above, Eq. (5), either fitted directly to the data or indirectly through Eq. (4), gives an empirical approximation to the total dependence of E_t upon $\langle n \rangle$

$$E_t = g(\langle n \rangle). \quad (6)$$

On the other hand, we have the total dependence of $\langle n \rangle$ upon E_t directly from the data

$$\langle n \rangle = f(E_t). \quad (7)$$

Clearly $g(\langle n \rangle)$ is an approximation to the inverse of $f(E_t)$, $g \approx f^{-1}$. Now we

construct the Arrhenius plot

$$F(\langle n \rangle) = F(f'(E_t)) \quad (8)$$

with

$$\langle n \rangle = f'(E_t) \propto \exp\left(\frac{-K}{\sqrt{E_t}}\right). \quad (9)$$

If the Arrhenius plot is linear it means the $f'(E_t) \equiv f$. Hence,

$$F(\langle n \rangle) = F(f(E_t)) = F(f(g(\langle n \rangle))) = F\left(f\left(f^{-1}(\langle n \rangle)\right)\right) = F(\langle n \rangle) \quad (10)$$

which is an identity, in the limit of $g = f^{-1}$.

The vicious circle of circular logic can be summarized as follows: given any function $y = y(x)$, substitute in it x with its value $x(y)$: $y = y(x(y))$. This gives always the identity $y = y$.

This result proves, if there was any doubt, that any function is completely correlated with its own inverse, and it is a great source of straight lines, irrespective of whether there is an autocorrelation or not.

The flaw, in our specific case, lies with the fact that Eq. (6) gives the *total empirical* dependence of E_t on $\langle n \rangle$ *irrespective of whether there is autocorrelation or not* and consequently *it is the approximate inverse of Eq. (7) and of Eq. (9)*. Hence the Uroboros, the snake eating its tail.

It is left for the diligent reader to test the universality of this method by applying the Uroboros to discover autocorrelations in the most venerable dependences in physics.

It is clear from the previous considerations that the authors' vicious circle does not prove anything, least of all autocorrelation.

In order to appreciate the true role of autocorrelations, let as before

$$\langle n \rangle = f(E_t) \quad (11)$$

and

$$E_t = E_t(E, \langle n \rangle) \quad (12)$$

where E is the excitation energy of the system. The dependence of $\langle n \rangle$ on E_t can be best explored differentially

$$\Delta \langle n \rangle = \frac{d \langle n \rangle}{d E_t} \Delta E_t. \quad (13)$$

Now, the change in transverse energy can be written

$$\Delta E_t = \left. \frac{\partial E_t}{\partial E} \right|_n \Delta E + \left. \frac{\partial E_t}{\partial \langle n \rangle} \right|_E \Delta \langle n \rangle \quad (14)$$

where the first term represents the change in E_t due to the change in excitation energy and the second term represents the trivial change in E_t due to autocorrelation. The partial derivative $\left. \frac{\partial E_t}{\partial \langle n \rangle} \right|_E$ at constant E (not the total derivative) tells us about the autocorrelation.

Substituting Eq. (14) in Eq. (13) we have

$$\Delta \langle n \rangle \left(1 - \frac{d \langle n \rangle}{d E_t} \left. \frac{\partial E_t}{\partial \langle n \rangle} \right|_E \right) = \frac{d \langle n \rangle}{d E_t} \left. \frac{\partial E_t}{\partial E} \right|_{\langle n \rangle} \Delta E \quad (15)$$

where the complete left hand side is the increase in $\langle n \rangle$ induced by ΔE corrected for the autocorrelation. Failure to appreciate the difference between $\left. \frac{\partial E_t}{\partial \langle n \rangle} \right|_E$ and $\frac{d E_t}{d \langle n \rangle}$ leads to disastrous circular reasoning.

In fact, if we write

$$\Delta E_t = \frac{d E_t}{d \langle n \rangle} \Delta \langle n \rangle \quad (16)$$

and substitute in Eq. (13) we obtain as the authors did

$$\Delta \langle n \rangle = \frac{d \langle n \rangle}{d E_t} \frac{d E_t}{d \langle n \rangle} \Delta \langle n \rangle = \Delta \langle n \rangle, \quad (17)$$

or the Uroboros (again).

Of course, the authors might argue that they know the first term of Eq. (14) to be close to zero because they believe the temperature changes at most by 1 MeV. Then indeed only the second term, or the autocorrelation would survive. This position is clearly untenable. In fact, were the authors correct, the derivative in Eq. (16) (which would now be the same as that in the second term of Eq. (14)) should have the value of approximately 20-40 MeV per

IMF, which is the measured mean transverse energy per IMF [16,17] for this particular system. Instead, it has the enormous value of 220 MeV per IMF! (See Fig. 1.) This alone should make the authors wonder and worry about constant temperature. In any case, as we pointed out above, their method will always give the same result, independent of the presence or absence of an autocorrelation.

For those who like to touch with their hands, here is a concrete example from an event generator code [3] which, *by construction*, generates complex fragment events according to

$$p = \exp(-B/T) \quad (18)$$

with

$$T = \sqrt{\frac{E}{a}}, \quad (19)$$

where T is the temperature and E is the excitation energy of the system. In other words, the linear Arrhenius plot is contained in the events by construction. By using the independent variable E instead of E_t , we are assured of avoiding the autocorrelation problem altogether.

We construct the Arrhenius plot $1/\langle n \rangle$ versus $1/\sqrt{E}$ which turns out to be linear as expected (circles in the bottom panel of Fig. 2). Then we notice that E and $\langle n \rangle$ can be plotted one against the other (circles in the top panel of Fig. 2) and we approximate

$$E = a + b \langle n \rangle \quad (20)$$

by fitting the dependence with a straight line. Then we substitute E with the value given by Eq. (20) and note, as the authors, that the Arrhenius plot constructed from this substitution (solid line in bottom panel of Fig. 2) remains linear over the range in E_t where Eq. (20) well represents the simulation. Hopefully by now the game has become rather tedious, and nobody would argue that the result is just due to an autocorrelation.

To verify the extent of autocorrelations in this model, we follow the same procedure as above but use E_t , like in the data. Now, we expect some autocorrelation. The question is: how much? We obtain a linear Arrhenius plot, whose slope should coincide with the slope of the previous plot after scaling E_t by $3/2$. (The model assumes $E_t = 2/3E$.) Indeed it does so to within better than 10 percent [3] (open squares of Fig. 2) *This difference in slope is the extent of the autocorrelation problem.* By redefining $\langle n \rangle$ through an increase




Fig. 2. Top panel: The excitation energy E as a function of the average IMF multiplicity from a binomial simulation [3] of the decay of a Xe nucleus (circles). The squares show the same correlation but using for E a scaled value of E_t , $E = 3/2E_t$. Bottom panel: The Arrhenius plot $1/\langle n \rangle$ versus $1/\sqrt{E}$ (circles) and $1/\langle n \rangle$ versus $1/\sqrt{3/2E_t}$ (squares). The solid line represents the (erroneous) “self-correlation” explanation proposed by the authors (see text).

of the element threshold (Z_{th}) in the definition of IMF (IMF: $Z_{\text{th}} \leq Z \leq 20$), we can substantially reduce even this small autocorrelation. Yet the authors would conclude that in both cases, using E or E_t , the linear Arrhenius plots are completely due to autocorrelation.

An even better procedure, which we can apply directly to experimental data, is the following. We can redefine the IMFs to be the fragments of a single element, say $Z=10$. Now the contribution of the $Z=10$ fragments to E_t is so small that it can be neglected and consequently autocorrelation is negligible as well. Because of the restriction of the definition of an IMF (one single Z), the probability of emitting an IMF becomes very small, and in fact, we observe that the distribution of multiplicities becomes Poissonian. The Arrhenius plots of $\log \langle n \rangle$ versus $1/\sqrt{E_t}$ remain linear and the linearity can be tested for each Z as in Fig. 3 without the worry of autocorrelation. The remarkable set of straight lines shown in Fig. 3, together with many other sets obtained for all the experiments we have analyzed, shows that thermal scaling is a *real*




Fig. 3. The average yield per event of the different indicated elements (symbols) as a function of $1/\sqrt{E_t}$. The lines are linear fits to the data.

physical feature quite safe from the dreaded fangs of the Uroboros snake. Yet the procedure employed by the authors will always yield an autocorrelation, regardless of whether one is present or not.

We now return to the first criticism, on which there is little to say.

The authors claim that there is no *a priori* reason for choosing a binomial instead of a constrained Poissonian, because they both fit the experimental multiplicity distributions equally well. We have no great argument with this. We actually like the Poissonian case and we would have chosen it if the data had not indicated a very high probability and a constraint (in size or temporal) to boot [3]. A binomial satisfied both requirements. By requiring a fixed number of tosses m , it incorporates both a scale and a constraint. There are other ways to achieve the same goal. For instance, we could chop the Poissonian above a prescribed n or do something else, like the authors did. In point of fact, when we restrict the definition of complex fragments to a single atomic number Z (see Fig. 3), the probability falls so much that the “binomials” become Poissonian on their own, removing us from the embarrassment of choice.

But binomials, Poissonians, and constrained Poissonians, all share the feature of being constructed upon *one* fragment probabilities. All of them construct the n -fragment probability P_n from the one fragment probability p . All are *reducible* and this is our experimental point! A Poissonian is *quintessentially*




Fig. 4. The empirical scaling of mZ_{th} for different Z_{th} (different symbols) for the reaction $^{36}\text{Ar}+^{197}\text{Au}$ at $E/A=110$ MeV.

reducible. Because of the loss of scale, its elementary probability is arbitrarily normalizable and coincides with the mean. Remembering that the Poissonian originates from the binomial, we get $\langle n \rangle = mp$ without however, being able to disentangle m from p . On the other hand, if we introduce a constraint, in whichever expression, we get back the approximate scale $m \approx Z_0 / \langle Z \rangle$ where Z_0 is the source size.

An example of a mass constraint in our binomial fits is given by the approximate scaling that we have observed [3] in the experiments:

$$m_{Z_{th}} Z_{th} \approx \text{constant} \quad (21)$$

shown in Fig. 4.

In view of the above, we do not see any merit to the authors criticism. Their function is reducible by construction, since it assumes *a priori* independent emission (see their appendix). Furthermore, their constraint reintroduces the scale, by the back door, that the Poissonian threw out the window. Reducibility is our key statement, binomial or Poissonian, as the case may be.

The authors seem to imply a big difference between the two, but the issue is reducibility. Clearly, our experimental claim that the n -fragment multiplicity

distributions are reducible to a single one fragment probability is not ever called into question by the authors.

In summary, the present preprint [9] attempts to give alternative or trivial explanations for the reducible and “thermal” nature of the experimental IMF excitation functions reported in [1–3]. We have demonstrated that the proposed “self-correlation” explanation for linear Arrhenius plots is based upon a flawed autocorrelation analysis involving circular reasoning. Equivalently, the authors have used a total derivative when they should have taken a partial derivative. This is a fatal mistake which produces the “self-correlation” result, independent of whether or not there are autocorrelations truly present.

References

- [1] L.G. Moretto *et al.*, Phys. Rev. Lett. **74**, 1530 (1995).
- [2] K. Tso *et al.*, Phys. Lett. B **361**, 25 (1995)
- [3] L.G. Moretto *et al.*, Phys. Rep. **287**, 249 (1997).
- [4] A.S. Botvina and D.H.E. Gross, Phys. Lett. B **344**, 6 (1995).
- [5] A. Del Zoppo *et al.*, Phys. Rev. Lett. **75**, 2288 (1995).
- [6] D. Gross, Phys. Rep. **279**, 120 (1997).
- [7] Alfredo Aranda, Jorge Alberto Lopez, and Zehua Wu, Phys. Rev. C **55**, 788 (1997).
- [8] J. Toke *et al.*, Rochester preprint UR-NSRL-423.
- [9] M.B. Tsang and P. Danielewicz, MSU preprint MSU-1069, and LANL e-print nucl-ex/9704010.
- [10] M.J. Huang *et al.*, Phys. Rev. Lett. **78**, 1648 (1997).
- [11] Z. Chen *et al.*, Phys. Lett. B **199**, 171 (1987).
- [12] D. Morrissey, W. Benenson, W. Friedman, Ann. Rev. Nucl. Part. Sci **44**, 27 (1994) and references therein.
- [13] F. Zhu *et al.*, Phys. Lett. B **282**, 299 (1992).
- [14] S. Albergo *et al.*, Nuovo Cimento A **89**, 1 (1985).
- [15] Y.G. Ma *et al.*, Phys. Lett. B **390**, 41 (1997).
- [16] R.T. de Souza *et al.*, Phys. Lett. B **300**, 29 (1993).
- [17] L. Phair, private communication.